



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68	A2	(11) International Publication Number: WO 99/05315 (43) International Publication Date: 4 February 1999 (04.02.99)
(21) International Application Number: PCT/GB98/02214 (22) International Filing Date: 24 July 1998 (24.07.98) (30) Priority Data: 9715942.0 28 July 1997 (28.07.97) GB 9727103.5 22 December 1997 (22.12.97) GB (71) Applicant (for all designated States except US): MEDICAL BIOSYSTEMS LTD. [GB/GB]; The Old Mill, Beaston Cross, Broadhempston, Nr. Totnes, Devon TQ9 6BX (GB). (72) Inventor; and (75) Inventor/Applicant (for US only): DENSHAM, Daniel, Henry [GB/GB]; The Old Mill, Beaston Cross, Broadhempston, Nr. Totnes, Devon TQ9 6BX (GB). (74) Agent: GILL JENNINGS & EVERY; Broadgate House, 7 Eldon Street, London EC2M 7LH (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: NUCLEIC ACID SEQUENCE ANALYSIS (57) Abstract The present invention relates to a method for determining the sequence of a polynucleotide, the method comprising the steps of: (i) reacting a target polynucleotide with a polymerase enzyme immobilised on a solid support, and the different nucleotides, under conditions sufficient for the polymerase reaction; and (ii) detecting the incorporation of a specific nucleotide complementary to the target polynucleotide, by measuring radiation.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Licchtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

NUCLEIC ACID SEQUENCE ANALYSIS

Field of the Invention

This invention relates to a method for determining the sequence of a polynucleotide.

5 Background of the Invention

The ability to determine the sequence of a polynucleotide is of great scientific importance. For example, the Human Genome Project is an ambitious international effort to map and sequence the three billion
10 basis of DNA encoded in the human genome. When complete, the resulting sequence database will be a tool of unparalleled power for biomedical research. The major obstacle to the successful completion of this project concerns the technology used in the sequencing process.

15 The principal method in general use for large-scale DNA sequencing is the chain termination method. This method was first developed by Sanger and Coulson (Sanger et al. Proc. Natl. Acad. Sci. USA 1977; 74: 5463-5467), and relies on the use of dideoxy derivatives of the four
20 nucleoside triphosphates which are incorporated into the nascent polynucleotide chain in a polymerase reaction. Upon incorporation, the dideoxy derivatives terminate the polymerase reaction and the products are then separated by gel electrophoresis and analysed to reveal the position at
25 which the particular dideoxy derivative was incorporated into the chain.

Although this method is widely used and produces reliable results, it is recognised that it is slow, labour-intensive and expensive.

30 An alternative sequencing method is proposed in EP-A-0471732, which uses spectroscopic means to detect the incorporation of a nucleotide into a nascent polynucleotide strand complementary to a target. The method relies on an immobilised complex of template and primer, which is
35 exposed to a flow containing only one of the different nucleotides. Spectroscopic techniques are then used to measure a time-dependent signal arising from the polymerase

5 catalysed growth of the template copy. The spectroscopic techniques described are surface plasmon resonance (SPR) spectroscopy, which measures changes in an analyte within an evanescent wave field, and fluorescence measuring techniques. However, limitations of this method are recognised; the most serious for the SPR technique being that, as the size of the copy strand grows, the absolute size of the signal also grows due to the movement of the strand out of the evanescent wave field, making it harder to detect increments. The fluorescence measuring techniques have the disadvantage of increasing background interference from the fluorophores incorporated on the growing nascent polynucleotide chain. As the chain grows, the background "noise" increases and the time required to detect each nucleotide incorporation needs to be increased. This severely restricts the use of the method for sequencing large polynucleotides.

20 There is therefore a need for an improved method for determining the sequence of polynucleotides which significantly increases the rate at which a polynucleotide is sequenced and which is preferably carried out by an automated process, reducing the complexity and cost associated with existing methods.

Summary of the Invention

25 The present invention is based on the realisation that the measurement of electromagnetic or other radiation can be used to detect a conformational and/or mass change in a polymerase enzyme which occurs when a nucleotide is incorporated into a nascent polynucleotide strand.

30 According to the present invention, a method for sequencing a polynucleotide comprises the steps of:

(i) reacting a target polynucleotide with a polymerase enzyme immobilised on a solid support, and the different nucleotides, under conditions sufficient for the polymerase reaction; and

35

(ii) detecting the incorporation of a specific nucleotide complementary to the target polynucleotide, by measuring radiation.

The radiation may be applied to a sample using a number of techniques, including surface-sensitive detection techniques, where a change in the optical response at a solid optical surface is used to indicate a binding interaction at the surface. In a preferred embodiment of the invention, the technique used is evanescent wave spectroscopy, in particular surface plasmon resonance (SPR) spectroscopy.

In an embodiment of the invention, the nucleotides used in the method include a blocking group at the 3' position, and optionally at the 5' position, which prevents incorporation of the nucleotides into the polynucleotide strand. However, the blocking groups may be selectively removed to allow incorporation to occur. By using the blocked nucleotides, it is possible for the method to be carried out using all the nucleotides present in the reaction at any one time. The selective removal of the blocking groups is carried out in a way that ensures the detection of each incorporated nucleotide. The method may therefore proceed on a "real-time" basis, to achieve a high rate of sequence analysis.

25 Description of the Drawings

The invention will be described by way of example only with reference to the following drawings, where:

Figure 1 is a schematic illustration of polynucleotide sequence analysis using SPR spectroscopy; and

30 Figure 2 illustrates the different response signals detected for the polymerisation of each of the different nucleotides.

Figure 3 illustrates the synthesis procedure for the double blocked nucleotides.

35 Description of the Invention

The present method for sequencing a polynucleotide involves the analysis of the kinetic interaction between a

polymerase enzyme, a target polynucleotide and a complementary nucleotide. Measurement of the kinetic interaction is carried out by monitoring the changes in or absorption of electromagnetic or other radiation that occurs if the reaction proceeds.

The term "polynucleotide" as used herein is to be interpreted broadly, and includes DNA and RNA, including modified DNA and RNA, as well as other hybridising nucleic acid-like molecules, e.g. peptide nucleic acid (PNA).

Typically, the method is carried out by applying electromagnetic radiation, by using the techniques of surface plasmon resonance or nuclear magnetic resonance. However, other techniques which measure changes in radiation may be considered, for example spectroscopy by total internal reflectance fluorescence (TIRF), attenuated total reflection (ATR), frustrated total reflection (FTR), Brewster angle reflectometry, scattered total internal reflection (STIR) or evanescent wave ellipsometry.

Techniques other than those requiring electromagnetic radiation are also envisaged, in particular photochemical techniques such as chemiluminescence, and gravimetric techniques including resonant systems such as surface acoustic wave (SAW) techniques and quartz crystal microbalance (QCM) techniques.

Surface plasmon resonance (SPR) spectroscopy is a preferred method, and measures the properties of a solution by detecting the differences in refractive index between the bulk phase of the solution and the evanescent wave region. Incident monochromatic light is reflected at a specific angle off a solid optical (sensor chip) surface on the opposite side to a sample under study. The light extends into the sample for a very short distance and is affected by an interaction at the surface.

Suitable sensor chips are known in the art. Typically, they comprise an optically transparent material, e.g. glass, and a thin reflective film, e.g. silver or gold. For a review of SPR spectroscopy see European Patent

Publication No. 0648328 (the entire disclosure of which is incorporated herein by reference).

Nuclear magnetic resonance (NMR) spectroscopy is another preferred method, and measures the magnetic properties of compounds. Nuclei of compounds are energetically orientated by a combination of applied magnetic field and radio-frequency radiation. When the energy exerted on a nucleus equals the energy difference between spin states (the difference between orientation parallel or anti-parallel to the direction of the applied fields), a condition known as resonance is achieved. The absorption and subsequent emission of energy associated with the change from one spin state to the other, is detected by a radio-frequency receiver.

An important aspect of the method of the present invention is the use of a polymerase enzyme immobilised onto a solid support. Immobilisation of the polymerase offers several important advantages for the success of this method. Firstly, the problem of random "noise" associated with measuring energy absorption in soluble molecules is reduced considerably. Secondly, the problem of noise from the interaction of any substrate (e.g. nucleotides) not directly involved with the polymerase is reduced, as the polymerase can be maintained within a specifically defined area relative to the field of measurement. This is particularly relevant if the technique used to measure the changes in radiation requires the measurement of fluorescence, as in TIRF, where background fluorescence increases as the nascent chain grows. Also, if SPR spectroscopy is used, the polymerase reactions are maintained within the evanescent wave field and so accurate measurements can be made irrespective of the size of the polynucleotide. Finally, as neither the target polynucleotide nor the oligonucleotide primer is irreversibly attached to the solid surface, it is relatively simple to regenerate the surface, to allow

further sequencing reactions to take place using the same immobilised polymerase.

Immobilisation may be carried out using standard procedures known in the art. In particular, immobilisation using standard amine coupling procedures may be used, with attachment of ligand-associated amines to, say, a dextran or N-hydroxysuccinimide ester-activated surface. In a preferred embodiment of the invention, the polymerase is immobilised onto a SPR sensor chip surface which maintains the polymerase in close proximity to the sensor surface where changes in the refractive index may be measured. Examples of procedures used to immobilise biomolecules to optical sensors are disclosed in EP-A-0589867, and Löfas et al., Biosens. Bioelectron. (1995) 10:813-822.

The polymerase used in the invention may be of any known type. For example, the polymerase may be any DNA-dependent DNA polymerase. If the target polynucleotide is a RNA molecule, then the polymerase may be a RNA-dependent DNA polymerase, i.e. reverse transcriptase, or a RNA-dependent RNA polymerase, i.e. RNA replicase. In a preferred embodiment of the invention, the polymerase is Taq polymerase. In a further preferred embodiment of the invention, the polymerase is either *E. coli* polymerase III holoenzyme (McHenry, Ann. Rev. Biochem. 1988; 57:519), T7 polymerase (Schwager et al., Methods in Molecular and Cellular Biology (1989/90); Vol.1(4):155-159, or Bacteriophage T7 gene 5 polymerase complexed with *E. coli* Thioredoxin (Tabor et al., J. Biol. Chem. (1987); 262:1612-1623). Each of these polymerase enzymes allows a binding with the target polynucleotide to occur with high fidelity and therefore maintains a polymerase-polynucleotide complex, even when polymerisation is not actively taking place.

The polymerase III holoenzyme is composed of three subassemblies that function to create the processive enzyme: (I) the polymerase core, including the polymerase subunit α ; (II) β -dimer subunit which acts as a bracelet-

like structure around DNA; and (III) a subassembly of two subunits, τ and γ , used to bind and hydrolyse ATP to form the β -dimer around the DNA.

As a first step in the sequencing process, the target
5 polynucleotide may be brought into contact with an
appropriate primer in hybridising/polymerisation buffer.
Typically, the buffer will be at a sufficiently high
temperature to disrupt (or melt) any secondary structures
that exist on the target polynucleotide. On cooling, the
10 primer will anneal to its complement on the target. This
sample may then be brought into contact with the
immobilised polymerase, to form the target polynucleotide/
polymerase complex.


In one embodiment of the invention, the addition of
15 the nucleotides is controlled so that the different
nucleotides are added sequentially to the polymerase/target
complex. For example, dGTP may be added and allowed to
flow over the polymerase/polynucleotide complex; any
incorporation is then detected. Unbound dGTP flows out of
20 the reaction site and a further nucleotide is introduced.
In this manner, the detection of a kinetic interaction can
be correlated to the particular nucleotide present at that
time and the polynucleotide sequence can therefore be
determined.

25 The method may also be carried out with all the
different nucleotides present. For this to be carried out
successfully, it is necessary for the nucleotides to
incorporate a blocking group at least at the 3' position,
but preferably at the 3' and 5' positions. The blocking
30 groups may be light-sensitive and can be removed by
applying light of a defined wave length, to release the
active molecule. If the nucleotides incorporate blocking
groups at both the 3' and 5' positions, the blocking groups
should be capable of being distinguished on the basis of
35 their spectral absorbancy, i.e. it should be possible to
remove selectively one of the blocking groups by applying
a specific wavelength of light which does not remove the

other blocking group. It is also desirable that the blocking group at the 3' position requires the light to be applied for a longer duration than that required to remove the blocking group at the 5' position. This allows the blocking groups to be distinguished by both spectral and temporal means.

Generally, the light-sensitive blocking groups undergo photolysis at wavelengths in the range from 200nm to 450nm. The blocking groups will typically be derived from a compound of the formula $R^1-[O-CO-]X$ wherein R^1 is a photolabile group and X is a leaving group. For example, R^1 is o-nitrobenzyl. Particularly preferred blocking groups include the o-nitrobenzyl protecting groups described in WO-A-92/10092 and WO-A-97/39151. These groups include nitroveratryloxycarbonyl (NVOC), nitropiperonyloxycarbonyl (NPOC), α -methyl-nitroveratryloxycarbonyl (MeNVOC), α -methyl-nitropiperonyloxycarbonyl (MeNPOC) and 1-pyrenylmethyloxycarbonyl (PYMOC).

A suitable 3' blocking group is a (4,5-dimethoxy-2-nitrobenzyl)oxycarbonyl group which can be formed by reaction of the nucleotide with a compound of formula (I):




wherein R is any suitable esterifying group, e.g. methyl. This blocking group can be selectively removed by a pulse of light with a wavelength of 360 nm.

A suitable blocking group at the 5' position is 1-(2-nitrophenyl)ethyl group (II):

5

10



15 wherein R is any suitable functional group, e.g. halogen. This blocking group may be selectively removed at a wavelength of 260 nm.

By way of example, double-blocked nucleotides are injected over the primed target polynucleotide (held in association with a high fidelity polymerase complex), and monochromatic light is focussed upstream of the polymerase at a wavelength sufficient to release the blocking group from the terminal phosphate of each nucleotide. The nucleotides are then able to flow over the bound polymerase, and incorporation into the nascent polynucleotide strand can occur. However, as the blocking group at the 3' position remains bound, only one nucleotide is incorporated. A measurement of the kinetic interaction will therefore provide information as to the particular nucleotide incorporated into the nascent chain. The polymerase used may be a high fidelity polymerase which does not dissociate readily from the target when the reaction stops. Alternatively, a competitive inhibitor may be used to prevent the polymerase dissociating from the target.

After measuring the incorporated nucleotide, a pulse of monochromatic light is focused on the blocking group

within the polymerase catalytic site, to remove the blocking group at the 3' position. The monochromatic light may be pulsed for a longer duration than that required for removal at the 5' position, and so only the blocking group associated with the nucleotide in the polymerase complex will undergo removal. This reduces the likelihood of the addition of nucleotides not associated with the polymerase complex.

Once the 3' blocking group is released, the polymerase reaction is allowed to continue as further nucleotides arrive at the polymerase reaction site. Uncontrolled polymerisation is prevented by alternating the pulses of light required to remove the blocking groups.

While it is preferred to use the double-blocked nucleotides, as described above, the procedure may also be carried out using nucleotides having a blocking group at the 3' position only. In this case, it is desirable to use a competitive inhibitor of the polymerase, which will reduce the probability of a nucleotide lacking a blocking group at the 3' position being incorporated into the nascent chain. A suitable competitive inhibitor of polymerase is carbonyldiphosphonate (COMDP).

The following Example illustrates the invention with reference to the drawings.

Example

The following analysis was carried out on a modified BIAcore 2000 system (BIAcore AB, UPPSALA, Sweden) with a sensor chip CM5 (Research grade, BIAcore AB) as the optical sensor surface. The instrument was provided with an integrated μ m-fluidic cartridge (IFC) which allows analysis in four cells by a single sample injection.

Preparation of polymerase

E. Coli polymerase III holoenzyme was prepared according to (Millard et al., Methods Enzymol. (1995); 262:22) using hydrophobic interaction chromatography on valyl-Sepharose, to purify the holoenzyme at high salt concentrations. After purification, the hollow enzyme was

concentrated using the ion-filtration technique described by Kirkegaard et al, Anal. Biochem. (1972); 50:122.

Immobilisation of the Polymerase

Immobilisation of the polymerase to the sensor chip surface was carried out according to (Jönsson et al., Biotechniques (1991); 11:620-627). Briefly, the sensor chip environment was equilibrated with Hepes buffer (10 mM Hepes, 150 mM NaCl, 0.05% surfactant P20 (BIAcore AB, Uppsala, Sweden), pH 7.4). Equal volumes of N-hydroxysuccinimide (0.1 M in water) and N-ethyl-n'-(dimethylaminopropyl) carbodiimide (EDC) (0.1 M in water) were mixed together and injected across the chip (CM5) surface, to activate the carboxymethylated dextran. The polymerase III Subassembly core (160 µl, 500 U) was mixed with 10 mM sodium acetate (100 µl, pH 5) and injected across the activated surface. Finally, residual N-hydroxysuccinimide esters on the sensor chip surface were reacted with ethanolamine (35 µl, 1 M in water, pH 8.5), and non-bound polymerase was washed from the surface. The immobilisation procedure was performed with a continuous flow of Hepes buffer (5 µl/min) at a temperature of 25°C.

Oligonucleotides

Two oligonucleotides were synthesised using standard phosphoramidite chemistry. The oligonucleotide defined as SEQ ID No. 1 was used as the target polynucleotide, and the oligonucleotide defined as SEQ ID No. 2 was used as the primer.

CAAGGAGAGGACGCTGTCTGTCGAAGGTAAGGAACGGACGAGAGAAGGGAGAG

30

SEQ ID No. 1

CTCTCCCTTCTCTCGTC

SEQ ID No. 2

The two oligonucleotides were reacted under hybridising conditions to form the target-primer complex.

The primed DNA was then suspended in buffer (20 mM Tris-HCl, pH 7.5, 8 mM MgCl₂, 4% (v/v) glycerol, 5 mM

dithiothreitol (DDT), 40 μ g bovine serum albumin) containing 21 μ g ssDNA binding-protein and the DNA pol III sub-assembly, required to form the bracelet-like structure (1.6 pmol β -dimer and 195 fmol γ subunits). 0.5 mM ATP was present together with 60 μ M carbonyldiphosphonate (COMDP). In this reaction, the γ subunit acts as a molecular matchmaker, hydrolysing ATP to place the β -dimer subunits onto the DNA to form the polymerase sub-assembly (Studwell et al, UCLA Symp. Mol. Cell. Biol. New Ser. 1990; 127: 153).

The primed DNA/sub-assembly complex was then injected over the polymerase III on the sensor chip surface at a flow-rate of 5 μ m/min, and allowed to bind to the polymerase via the action of the γ subunits.

In this experiment, magnesium and ATP are needed for the Pol III to bind to the primed DNA. However, magnesium also promotes removal of the primer by the proof-reading 3' - 5' exonuclease activity. This problem is circumvented by including the carbonyldiphosphonate, which is a competitive inhibitor of polymerase activity (a pol III lacking the 3'-5' exonuclease activity may be used to avoid this particular problem).

A continuous flow of 60 μ M carbonyldiphosphonate was maintained over the chip surface, to prevent the exonuclease activity from removing the primer from the target DNA.

Nucleotides Incorporating Two Blocking Groups

Each nucleotide (dCTP, dTTP, dGTP and dATP) contained a 1-(2-nitrophenyl)ethyl blocking group at the 5' position, and a (4,5-dimethoxy-2-nitrobenzyl) oxycarbonyl blocking group at the 3' position as shown in Figure 3. The synthesis of the double blocked nucleotides was as follows:
Stage 1:- Synthesis of (4,5-dimethoxy-nitrobenzyl)oxycarbonyl-nucleoside triphosphate.

The same overall method was applied to dGTP, dCTP and dTTP. A mixture of dATP dihydrate (0.4mmol) and approximately 3mmol of 4,5-dimethoxy-2-

nitrophenyldiazomethane, freshly prepared from 900mg (4 mmol) of 4,5-dimethoxy-2-nitrophenylhydrazine (synthesized by treatment of 6-nitroveraldehyde with hydrazine monohydrate in chloroform by the procedure of Wootton and
5 Trentham, Photochemical Probes in Biochemistry (Nielsen, P.E., Ed,) NATO ASI Ser. C, Vol. 272, p277-296 (1989), was stirred in 15 ml of DMSO at room temperature in the dark for 40 h. Monitoring of the reaction by TLC in a
10 chloroform/methanol (5:1 v/v) solvent system revealed the appearance of a spot with Rf 0.54 corresponding to the caged nucleotide. DMSO, unreacted diazo compound, and reaction products with low polarity were removed by
15 repetitive extraction with 60 ml of ether. The residual material, which, among other substances, contained unreacted nucleotide and the desired product, was dissolved in a minimal amount of chloroform and separated by flash
20 chromatography on a silica column (3 x 30 cm). Elution using 100% chloroform and methanol/chloroform (95:5 v/v) removed the hydrophobic side products of 4,5-dimethoxy-2-nitrophenyldiazomethane from the column. The fractions
25 were dried on a rotary evaporator. 78 mg of the caged product was then lyophilised. The overall yield was 45%. The 3' blocked 4,5-dimethoxy-2-nitrobenzyl oxycarbonyl dATP was isolated directly with higher purity by
30 preparative reverse-phased HPLC from the crude product.
Stage 2:- Addition of the 5' 1-(2-nitrophenyl)ethyl group to the 3' 4,5-dimethoxy-2-nitrobenzyl oxycarbonyl blocked dATP.

A mixture of 4,5-dimethoxy-2-nitrobenzyl oxycarbonyl
30 5' dATP (0.4mmol) and approximately 3mmol of 1-(2-nitrophenyl)diazoethane, freshly prepared from 716.7 mg (4mmol) of hydrazone of 2-nitroacetophenone (synthesized by treatment of 2-nitroacetophenone with hydrazine monohydrate in ethanol) and 2.9g (30mmol) of MnO₂ (90%) in 20 ml of
35 chloroform by the procedure of walker et al (walker et al, Methods Enzymol. 1989; 172:288-301), was stirred in 15 ml of DMSO at room temperature in the dark for 40 h.

Monitoring the reaction by TLC in a chloroform/methanol (5:1 v/v) solvent system revealed the appearance of a pair of spots with Rf 0.68 and Rf 0.58, corresponding to the two diastereoisomers of the axial and the two diastereoisomers of the equatorial form of the 1(2-nitrophenyl)ethyl ester of 4,5-dimethoxy-2-nitrobenzyl oxycarbonyl 5' dATP, respectively. DMSO, unreacted diazo compound, and reaction products with low polarity were removed by repetitive extraction with 50 ml ether.

The residual material, which contained among other substances unreacted 4,5-dimethoxy-2-nitrobenzyl oxycarbonyl 5' dATP and the desired double blocked dATP, was dissolved in a minimal amount of chloroform and separated by flash chromatography on a silica column 3 x 30 cm. Elution using 100% chloroform removed hydrophobic side products of 1-(2-nitrophenyl)diazoethane from the column. The product was dried on a rotary evaporator. Lyophilization gave 74 mg of the caged compound. The overall yield was 57%.

0.2 mM of each nucleotide was present in the polymerisation buffer (1 mM Tris-HCl pH 8.8, 5 mM KCl, 0.15 mM MgCl₂, 0.01 % (w/v) gelatin).

DNA Sequencing

Figure 1 shows a SPR sensing system and fluidic cell (7), having a means for applying electromagnetic radiation (1) to a sensor chip (2) with an immobilised polymerase enzyme (3) at the sensor surface, an inlet (4) for introducing the different nucleotides into the cell and two focusing assemblies (5) and (6) for pulsing monochromatic light into the cell.

The different nucleotides are introduced into the fluidic cell (7) at a flow rate of 30 µl/min., at a temperature of 25°C and a data collection rate of 10 Hz. As the nucleotides pass the focusing assembly (5), monochromatic light at a wavelength of 260 nm is pulsed to remove the blocking group at the 5' position. The nucleotides then flow over the sensor chip (2) and contact

the target polynucleotide/ polymerase complex (3) which is held in place by the β -dimer sub-assembly. Since the 3' position on the primer sequence is free to react, polymerisation may take place as a nucleotide is incorporated onto its complement on the target polynucleotide. This incorporation is then detected by the monochromatic p-polarised light of the SPR device. No further polymerisation occurs, since the incorporated nucleotide has a blocking group at the 3' position. Monochromatic light of wavelength 360 nm is then pulsed by the focusing assembly (6) at the site of polymerisation. The high flow rate in the fluidic cell ensures that nucleotides not bound to the polymerase are removed from the cell before sufficient energy has been absorbed to release their 3' blocking groups.

Once the 3' blocking group has been released from the polymerised nucleotide, further polymerisation may occur.

Figure 2 shows the results from the sequencing experiment with each nucleotide incorporated into the nascent chain being detected. The results show a sequence complementary to that of SEQ ID No. 1.

CLAIMS

1. A method for sequencing a polynucleotide, comprising the steps of:

5 (i) reacting a target polynucleotide with a polymerase enzyme immobilised on a solid support, and the different nucleotides, under conditions sufficient for the polymerase reaction; and

10 (ii) detecting an effect consequent on the incorporation of a specific nucleotide complementary to the target polynucleotide.

2. A method according to claim 1, wherein the effect in step (ii) is detected by measuring radiation.

15 3. A method according to claim 1 or claim 2, wherein steps (i) and (ii) are conducted with each of the different nucleotides in turn, until incorporation is detected, and then repeated.

4. A method according to claim 1 or claim 2, wherein step (i) is conducted with all the nucleotides present.

20 5. A method according to any preceding claim, wherein the nucleotides comprise a 3' blocking group which is removed after the polymerase reaction.

25 6. A method according to claim 5, wherein the blocking group can be selectively removed by pulsed monochromatic light.

30 7. A method according to claim 5 or claim 6, wherein the nucleotides comprise a further blocking group at the terminal phosphate group of the triphosphate chain, and the further blocking group is removed prior to the removal of the 3' blocking group.

8. A method according to claim 7, wherein the further blocking group can be selectively removed by pulsed monochromatic light under conditions different from those required to remove the 3' blocking group.

35 9. A method according to claim 8, wherein the further blocking group is removed by pulsing the monochromatic

light for a duration different from that required to remove the 3' blocking group.

10. A method according to any preceding claim, wherein step (i) further comprises introducing a competitive
5 inhibitor of the polymerase enzyme.

11. A method according to any preceding claim, wherein the target polynucleotide of step (i) is bound to the polymerase enzyme by a β_2 dimer complex.

12. A method according to any preceding claim, wherein the
10 polymerase is *E. coli* DNA polymerase III or T7 polymerase.

13. A method according to any of claims 1 to 11, wherein the polymerase is Taq polymerase.

14. A method according to any of claims 1 to 11, wherein the polymerase is reverse transcriptase.

15. 15. A method according to any preceding claim, wherein step (ii) comprises detection of a change in resonance signal over time.

16. A method according to any preceding claim, wherein the radiation is electromagnetic.

20 17. A method according to claim 16, wherein the electromagnetic radiation is in the infra-red spectrum.

18. A method according to any preceding claim, wherein step (ii) comprises using surface plasmon resonance.

25 19. A method according to claim 16, wherein the electromagnetic radiation is in the radio-frequency spectrum.


20. A method according to claim 19, wherein the incorporation of a nucleotide is detected using NMR.

30 21. A method according to any preceding claim, wherein the polynucleotide is DNA.

22. A sensor chip comprising a polymerase enzyme immobilised thereon.

35 23. A nucleotide comprising a blocking group at the 3' position and at the terminal phosphate group of the triphosphate chain, wherein the two blocking groups are removable by monochromatic light of different wavelengths.

24. A nucleotide according to claim 23, wherein the blocking groups are derived from a compound of the formula



5

wherein R^1 is a photolabile group and X is a leaving group.

25. A nucleotide according to claim 23 or claim 24, wherein the blocking group at the 3' position is an o-nitrobenzyloxycarbonyl group.

10 26. A nucleotide according to any of claims 23 to 25, wherein the blocking group at the terminal phosphate is an o-nitrobenzyl group.

15 27. A nucleotide according to any of claims 23 to 26, wherein the blocking group at the 3' position is a (4,5-dimethoxy-2-nitrobenzyl)oxycarbonyl group.

28. A nucleotide according to any of claims 23 to 27, wherein the blocking group at the terminal phosphate is a 1-(2-nitrophenyl)ethyl group.

20 29. An apparatus for sequencing a polynucleotide, comprising an optical sensor chip, a light source, an imaging device and a photodetector, wherein the sensor chip is as defined in claim 22.

1/5




Fig. 1

2/5




Fig. 2a




Fig. 2b

4/5




Fig. 2c

5 / 5




Fig. 3

THIS PAGE BLANK (USPTO)

(51) International Patent Classification ⁶ : C12Q 1/68		A3	(11) International Publication Number: WO 99/05315
			(43) International Publication Date: 4 February 1999 (04.02.99)
(21) International Application Number: PCT/GB98/02214 (22) International Filing Date: 24 July 1998 (24.07.98) (30) Priority Data: 9715942.0 28 July 1997 (28.07.97) GB 9727103.5 22 December 1997 (22.12.97) GB (71) Applicant (for all designated States except US): MEDICAL BIOSYSTEMS LTD. [GB/GB]; The Old Mill, Beaston Cross, Broadhempston, Nr. Totnes, Devon TQ9 6BX (GB). (72) Inventor; and (75) Inventor/Applicant (for US only): DENSHAM, Daniel, Henry [GB/GB]; The Old Mill, Beaston Cross, Broadhempston, Nr. Totnes, Devon TQ9 6BX (GB). (74) Agent: GILL JENNINGS & EVERY; Broadgate House, 7 Eldon Street, London EC2M 7LH (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 22 April 1999 (22.04.99)	
(54) Title: NUCLEIC ACID SEQUENCE ANALYSIS			
(57) Abstract			
<p>The present invention relates to a method for determining the sequence of a polynucleotide, the method comprising the steps of: (i) reacting a target polynucleotide with a polymerase enzyme immobilised on a solid support, and the different nucleotides, under conditions sufficient for the polymerase reaction; and (ii) detecting the incorporation of a specific nucleotide complementary to the target polynucleotide, by measuring radiation.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No

PC1/GB 98/02214

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 90 13666 A (AMERSHAM INT PLC) 15 November 1990 see the whole document ---	1-22,29
A	WO 93 21340 A (MEDICAL RES COUNCIL ;BRENNER SYDNEY (GB); ROSENTHAL ANDRE (GB)) 28 October 1993 see the whole document ---	1-22,29
A	WO 90 05303 A (PHARMACIA AB) 17 May 1990 see the whole document ---	1-22,29
A	WO 91 06678 A (STANFORD RES INST INT ;TSIEN ROGER Y (US)) 16 May 1991 see the whole document ---	1-22,29
-/--		

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

4 November 1998

Date of mailing of the international search report

25.02.1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hagenmaier, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB98/02214

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SCHWARZ T: "DETECTION OF NUCLEIC ACID HYBRIDIZATION USING SURFACE PLASMON RESONANCE" TRENDS IN BIOTECHNOLOGY, vol. 9, no. 10, 1 October 1991, pages 339-340, XP000247400 see the whole document ---	1-22,29
A	WO 94 21822 A (KOESTER HUBERT) 29 September 1994 see the whole document -----	1-22,29

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB 98/ 02214

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see FURTHER INFORMATION sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-22, 29 (completely)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-22, 29 (completely)

Method for sequencing a polynucleotide, comprising the steps of (i) reacting a target polynucleotide with a polymerase enzyme immobilised on a solid support, and the different nucleotides, under conditions sufficient for the polymerase reaction; and (ii) detecting the incorporation of a specific nucleotide complementary to the target polynucleotide, by measuring radiation as well as a sensor chip comprising a polymerase enzyme immobilised thereon and an apparatus comprising that sensor chip, a light source, an imaging device and a photodetector.

2. Claims: 23-28 (completely)

A nucleotide comprising a blocking group at the 3' position and at the terminal phosphate group of the triphosphate chain, wherein the two blocking groups are removable by monochromatic light of different wavelengths.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/02214

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9013666 A	15-11-1990	CA 2045505 A	12-11-1990
		EP 0471732 A	26-02-1992
		JP 4505251 T	17-09-1992

WO 9321340 A	28-10-1993	AT 159766 T	15-11-1997
		AU 4020893 A	18-11-1993
		CA 2133956 A	28-10-1993
		DE 69314951 D	04-12-1997
		DE 69314951 T	19-03-1998
		EP 0640146 A	01-03-1995
		ES 2110604 T	16-02-1998
		JP 7507681 T	31-08-1995

WO 9005303 A	17-05-1990	SE 462454 B	25-06-1990
		AT 136651 T	15-04-1996
		DE 68926255 D	15-05-1996
		DE 68926255 T	31-10-1996
		EP 0589867 A	06-04-1994
		JP 2815120 B	27-10-1998
		JP 4501605 T	19-03-1992
		SE 8804073 A	10-11-1988
		US 5436161 A	25-07-1995
		US 5242828 A	07-09-1993

WO 9106678 A	16-05-1991	CA 2044616 A	27-04-1991
		EP 0450060 A	09-10-1991

WO 9421822 A	29-09-1994	AU 687801 B	05-03-1998
		AU 6411694 A	11-10-1994
		CA 2158642 A	29-09-1994
		EP 0689610 A	03-01-1996
		JP 8507926 T	27-08-1996
		US 5622824 A	22-04-1997
		US 5851765 A	22-12-1998

THIS PAGE BLANK (USPTO)